



저작자표시-비영리-변경금지 2.0 대한민국

이용자는 아래의 조건을 따르는 경우에 한하여 자유롭게

- 이 저작물을 복제, 배포, 전송, 전시, 공연 및 방송할 수 있습니다.

다음과 같은 조건을 따라야 합니다:



저작자표시. 귀하는 원저작자를 표시하여야 합니다.



비영리. 귀하는 이 저작물을 영리 목적으로 이용할 수 없습니다.



변경금지. 귀하는 이 저작물을 개작, 변형 또는 가공할 수 없습니다.

- 귀하는, 이 저작물의 재이용이나 배포의 경우, 이 저작물에 적용된 이용허락조건을 명확하게 나타내어야 합니다.
- 저작권자로부터 별도의 허가를 받으면 이러한 조건들은 적용되지 않습니다.

저작권법에 따른 이용자의 권리는 위의 내용에 의하여 영향을 받지 않습니다.

이것은 [이용허락규약\(Legal Code\)](#)을 이해하기 쉽게 요약한 것입니다.

[Disclaimer](#)

의학석사 학위논문

The Prevalence of Vitamin D Deficiency
in Patients Undergoing Total Knee Arthroplasty
: A Propensity Score Matching Analysis

슬관절 인공관절 전치환술을 받은 환자에서
비타민 디 결핍의 유병률
: 성향 점수 매칭 분석

울산대학교 대학원

의학과

박기봉

The Prevalence of Vitamin D Deficiency
in Patients Undergoing Total Knee Arthroplasty
: A Propensity Score Matching Analysis

지 도 교 수 빈 성 일

이 논문을 의학석사 학위 논문으로 제출함

2022년 2월

울 산 대 학 교 대 학 원

의 학 과

박 기 봉

박기봉의 의학석사 학위 논문을 인준함

심사위원 김종민 (인)

심사위원 빈성일 (인)

심사위원 이범식 (인)

울 산 대 학 교 대 학 원

2022년 2월

Abstract (English)

Purpose: We investigated and compared the serum 25-OH vitamin D [25(OH)D] level and prevalence of vitamin D deficiency (VDD) between patients who underwent total knee arthroplasty (TKA) and healthy participants through a matched analysis.

Methods: The case group consisted of 824 patients who underwent TKA and the control group -2,794 healthy participants examined at our institution. The control group was adjusted for sex, age, weight, body mass index (BMI), blood chemistry, and season of sampling through propensity score matching (PSM). After PSM, 501 and 721 patients were matched to the case and control group, respectively. We classified the patient who had BMI more than 30 kg/m² as within the obesity range. Levels of blood chemistry including 25(OH)D were examined and VDD was defined as <20 ng/mL.

Results: After PSM, the serum 25(OH)D levels of both groups were significantly different, at 15.3 and 19.9 ng/mL, respectively, and the VDD prevalence in each group was 75.0% and 59.4%, respectively. In the case group, no differences were observed according to sex and obesity status. The seasonal VDD prevalence was significantly different between winter and autumn in the case group. The difference in seasonal VDD prevalence between the two groups was greatest in summer (24.1% points).

Conclusions: The patients who underwent TKA had lower serum 25(OH)D level and higher VDD prevalence compared to the healthy participants who matched using PSM. There were no differences in VDD rates by sex or obesity and the VDD prevalence was more than 70% during all season. Therefore, in patients undergoing TKA, general attention to VDD is required regardless of sex, obesity, and season.

Level of Evidence: Retrospective cohort study; Level III

Keywords: 25-OH vitamin D, knee, osteoarthritis, total knee arthroplasty, health checkup, vitamin D deficiency

Index

| | |
|--------------------------------|-----|
| Abstract (English)..... | i |
| Index for table & figures..... | iii |
| Introduction..... | 1 |
| Materials and Methods..... | 2 |
| Results..... | 5 |
| Discussion..... | 13 |
| Conclusion..... | 16 |
| Reference..... | 17 |
| 요약 (국문)..... | 19 |

Index for table & figures

| | |
|--|----|
| Table 1. Results of balance test..... | 6 |
| Table 2. Vitamin D status according to season of sampling in osteoarthritis (OA) and control group..... | 11 |
| Figure 1. The balance of covariates in the all dataset and matched set was evaluated by measuring standardized differences in means. All standardized mean differences in the baseline variables were <0.20, except for age (asterisk)..... | 7 |
| Figure 2. Vitamin D status in both group (OA and control group). The prevalence of vitamin D deficiency, insufficiency and sufficiency in the OA group was 74.8%, 18.4%, and 6.8%, respectively. The prevalence of vitamin D deficiency, insufficiency, and sufficiency in the control group was 59.5%, 24.5%, and 16.0%, respectively..... | 8 |
| Figure 3. The percentage of vitamin D deficiency (VDD) according to sex in osteoarthritis (OA) and control group. The percentage of VDD in the OA group was significantly higher than in the control group for both sexes. There was no significant difference between female and male in the OA group (p=0.06), and in the control group, the percentage of VDD in females (63.6%) was significantly higher than that of the males (47.6%). Asterisk (*) indicates p-value <0.01..... | 9 |
| Figure 4. The percentage of vitamin D deficiency (VDD) according to obesity in osteoarthritis (OA) and control group. There was no significant difference in the percentage of VDD in the OA and control groups in obesity patients. In non-obesity patients, there was significant difference between the two groups..... | 10 |
| Figure 5. The percentage of vitamin D deficiency according to the season of sampling in the osteoarthritis (OA) and control group. Between the two groups, a significant difference was observed in the percentage of VDD in summer, autumn, and winter (asterisk). Seasonal differences in VDD prevalence in the OA group were significant only in winter and autumn. In the control group, there were significant differences in VDD prevalence between summer and spring and between winter and summer..... | 12 |

Introduction

Vitamin D plays significant roles in the musculoskeletal system such as on cartilage health, muscle strength, sarcopenia, and fall prevention in the elderly.¹⁻³ Vitamin D status associated with the postoperative outcomes and rehabilitation of patients undergoing various orthopedic surgeries,⁴⁻⁶ and patients with vitamin D deficiency (VDD) undergoing total knee arthroplasty (TKA) have significantly worse outcomes.⁷⁻⁹

High rates of VDD have been demonstrated in various patient populations, including patients undergoing total joint arthroplasties.¹⁰ Maier et al.¹¹ have reported that over 60% of patients had VDD in a study of 1,083 patients aged >70 years and admitted to an orthopedic surgery department. Goula et al.¹² have evaluated the serum 25-OH vitamin D [25(OH) D] status in patients with knee or hip osteoarthritis (OA) scheduled for joint replacement and have reported that 81.7% of patients had VDD. Regarding the role of 25(OH)D level in TKA, Jansen and Haddad⁸ have reported a high prevalence of VDD in elderly patients with advanced knee OA scheduled for TKA, and Piuzzi et al.¹⁰ have reported that 26.9% of patients had VDD.

In a study using the National Health and Nutrition Examination Survey (NHANES) data collected from 2001 to 2004, the mean serum 25(OH)D level was 24 ng/mL and the prevalence of VDD was 77%.¹³ In a study using the fourth Korea NHANES conducted in 2008, VDD was found in 47.3% of men and 64.5% of women.¹⁴

To the best of our knowledge, no study has yet examined serum 25(OH)D levels and compared the prevalence of VDD in patients undergoing TKA and matched healthy individuals. Therefore, we performed a matched analysis of patients who underwent TKA and healthy individuals who were examined and compared the serum 25(OH)D level and VDD in the two groups. We hypothesized that there would be a significant difference in serum 25(OH)D levels or in the prevalence of VDD between the two groups in matched analysis.

Materials and Methods

This study was approved by our institutional review board (UUH-2021-05-019), and informed consent was obtained from all study participants.

Study design and setting

This was as a retrospective single-center study with a matched case-control design to address the purpose of this study. All study participants lived in a city located at 35.6 degrees latitude and were of a single Asian ethnicity.

Patients

We performed a retrospective review of the medical records from January 2012 to December 2020. Based on the records, 898 patients underwent primary TKA. Th patients recruited underwent TKA following presentation with moderate or advanced OA (Kellgren-Lawrence [15] grade III-IV, K-L grade III-IV) of the knee. First, patients with metabolic diseases such as hyperparathyroidism and Paget's disease were excluded. Second, patients with serum 25(OH)D levels <3 mg/mL and >70 mg/mL were excluded. Then, the late laboratory findings of patients who underwent staged bilateral surgery were excluded because the serum 25(OH)D levels could be changed after the first surgery among staged bilateral TKA. Therefore, 824 patients were included in the case group (OA group). Meanwhile, the criteria for inclusion in the control group were at least 55 years of age, no history of TKA, and had undergone regular health checkup in our institution. Altogether, 2,794 participants were included in the control group.

Propensity score matching strategy

The basic principle of propensity score matching (PSM) analysis was to estimate the propensity score (PS) for all observations in the case group and to match it with the observations in the control group with the closest PS value. Moreover, selection bias in selecting case and control groups can be reduced through PSM analysis, and any effect can be compared more accurately by controlling the influence of variables that can affect the results. The purpose of PSM analysis was to balance the two groups, not the number of observations in the two groups. Therefore, if there was no observation with the closest PS value in the control group, matching would not be possible. The reason why the number of observations was lost in the case group was the same with the reason as above. To reduce the bias and potential confounding among patients in the two groups, we performed careful adjustment for different distributions of baseline characteristics using a 2:1 PSM analysis with nearest-neighbor

method based on a greedy matching algorithm that sorts by estimated PS. PSM was conducted by a 2:1 nearest-neighbor matching using a caliper size of 0.2. The balance of covariates in the matched groups was evaluated by measuring their standardized differences in means. All standardized mean differences in the baseline variables were <0.20 (20%). In the propensity-score-matched pairs, the risks of each outcome were compared by logistic regression using generalized estimating equations for categorical variables or by the linear mixed model for continuous variables that accounted for the clustering of matched pairs. We matched the groups according to sex, age, weight, body mass index (BMI), blood chemistry, and season of sampling. The blood chemistry included measurements of serum levels of alkaline phosphatase, calcium, and phosphorus, which are known to be associated with vitamin D level. The blood chemistry was a routine test item to evaluate a patient's health condition before TKA; hence, it was planned to be used as a PSM analysis item. The program performs all analyses using the R software (version 4.0.2).

Categorization of vitamin D level

The serum 25(OH)D levels were measured in routine blood samples taken from patients at their pre-admission visit using radioimmunoassay and checked for vitamin D status. The patients were categorized into three groups according to their vitamin D status. VDD, insufficiency, and sufficiency were defined as a serum 25(OH)D level of <20 ng/mL, 20–29 ng/mL, and 30–150 ng/mL, respectively.¹⁶ The distribution of vitamin D categories was checked in the two groups, and the vitamin D levels were compared by category in both groups.

Obesity

BMI is a person's weight in kilograms divided by the square of height in meters. If the patient's BMI was ≥ 30.0 , we classified the patient as within the obesity range.¹⁷

Season of sampling

The season of sampling was divided into spring (March, April, and May), summer (June, July, and August), autumn (September, October, and November), and winter (December, January, and February).

Statistical analysis

Data are presented as mean with ranges. The Chi-square test was performed to verify the difference in the ratio of vitamin D categories between both groups. A p-value of <0.05 was considered significant. Statistical analyses

were performed using the SPSS version 24 (SPSS Inc., Chicago, IL, USA).

Results

Demographics


Before matching, the demographics of the two groups showed significant differences in age, sex, BMI, alkaline phosphatase, and phosphorus (Table 1). The average age of the 3,618 patients was 63.7 (range, 55–90) years; 1,908 (52.7%) were women and 1,710 (47.3%) were men. The average weight was 63.5 (range, 32.3–103.4) kg, and the average BMI was 24.6 (range, 14.5–45.3) kg/m². Here, a total of 1,222 patients were matched 2:1 through PSM, and no significant difference was found in all variables between the two groups, except for age (Figure 1). The average age of the 1,222 patients was 67.3 (range, 55–90) years; 939 (76.8%) patients were women and 283 (23.1%) were men. The average weight was 61.6 (range, 34.8–102.0) kg and the average BMI was 25.3 (range, 14.9–40.2) kg/m².

Table 1. Results of balance test

| | | Baseline characteristics of all patients | | | | Baseline characteristics of propensity score matched patients | | | |
|--------|--------|--|-------------------------------|-------|-------------|---|-----------------------------|------------|-------------|
| | | OA group (n=824) | Control group (n=2,794) | SMD | p- value | OA group (n=501) | Control group (n=721) | SMD | p- value |
| | | <i>n</i> (%) | <i>n</i> (%) | | | <i>n</i> (%) | <i>n</i> (%) | | |
| Sex | Female | 721 (87.5) | 1,187 (42.5) | -1.36 | 0.00 | 403 (80.4) | 98 (19.6) | -0.15 | 0.01 |
| | Male | 103 (12.5) | 1,607 (57.5) | | | 98 (19.6) | 185 (25.7) | | |
| Season | Spring | 218 (26.5) | 715 (25.6) | 0.01 | 0.97 | 126 (25.1) | 188 (26.1) | -0.01 | 0.89 |
| | Summer | 198 (24.0) | 683 (24.4) | | | 131 (26.1) | 182 (25.2) | | |
| | Autumn | 191 (23.2) | 658 (23.6) | | | 107 (21.4) | 155 (21.5) | | |
| | Winter | 217 (26.3) | 738 (26.4) | | | 137 (27.3) | 196 (27.2) | | |
| | | Mean±SD | Mean±SD | SMD | p- value | Mean±SD | Mean±SD | SMD | p- value |
| Age | | 70.1±6.1 | 61.8±5.8 | -1.37 | 0.00 | 68.1±5.5 | 66.7±6.5 | -0.26 | 0.00 |
| Weight | | 62.2±9.8 | 63.8±10.4 | 0.16 | 0.00 | 61.4±9.4 | 61.8±10.4 | 0.04 | 0.55 |
| BMI | | 26.6±3.5 | 24.0±2.9 | -0.75 | 0.00 | 25.6±2.9 | 25.2±3.3 | -0.13 | 0.03 |
| ALP | | 73.3±23.8 | 67.8±20.7 | -0.23 | 0.00 | 71.5±22.8 | 71.9±25.6 | 0.02 | 0.08 |
| Ca | | 9.3±0.5 | 9.3±0.4 | 0.12 | 0.00 | 9.3±0.5 | 9.3±0.4 | - 0.002 | 0.95 |
| P | | 3.61±0.56 | 3.45±0.53 | -0.29 | 0.00 | 3.6±0.58 | 3.6±0.5 | -0.03 | 0.58 |

OA, osteoarthritis; SMD, standardized mean difference; SD, standard deviation; BMI, body mass index; ALP, alkaline phosphatase; Ca, calcium; P, phosphorus


Figure 1. The balance of covariates in the all dataset and matched set was evaluated by measuring standardized differences in means. All standardized mean differences in the baseline variables were <0.20, except for age (asterisk)



Vitamin D status

The serum 25(OH)D levels of the patients were normally distributed, with a mean value of 19.1 (range, 3–66.6) ng/mL. The average serum 25(OH)D level was significantly lower in the OA group (15.3 ng/mL) than that in the control group (19.9 ng/mL, $p < 0.001$). When categorized using a 20 ng/mL cutoff, the VDD prevalence was 75.0% and 59.4% in the OA and control group, respectively (Figure 2). The prevalence of vitamin D insufficiency was 18.4% and 24.5% in the OA and control group, respectively. The prevalence of vitamin D sufficiency was 6.8% and 15.9% in the OA and control group, respectively ($p < 0.001$).


Figure 2. Vitamin D status in both group (OA and control group). The prevalence of vitamin D deficiency, insufficiency and sufficiency in the OA group was 74.8%, 18.4%, and 6.8%, respectively. The prevalence of vitamin D deficiency, insufficiency, and sufficiency in the control group was 59.5%, 24.5%, and 16.0%, respectively.



VDD according to sex

The VDD percentage according to sex is shown in Figure 3. As a result of comparison between the same sexes, the percentage of VDD in the OA group was significantly higher than that in the control group for both sexes (female, $p < 0.01$; male, $p < 0.01$). As a result of comparing men and women in the same group, there was no significant difference between men and women in the OA group ($p = 0.06$), and in the control group, the percentage of VDD in women (63.6%) was significantly higher than that of men (47.6%).


Figure 3. The percentage of vitamin D deficiency (VDD) according to sex in osteoarthritis (OA) and control group. The percentage of VDD in the OA group was significantly higher than in the control group for both sexes. There was no significant difference between female and male in the OA group ($p = 0.06$), and in the control group, the percentage of VDD in females (63.6%) was significantly higher than that of the males (47.6%). Asterisk (*) indicates p -value < 0.01 .



VDD according to obesity

The VDD percentage according to patients' obesity status is shown in Figure 4. No significant difference was noted in the percentage of VDD in the OA group (75.8%) and control group (67.2%) in obesity patients ($p=0.39$). However, in non-obesity patients, 74.7% (349/467) of the OA group had VDD, whereas 58.7% (387/659) of the control group had VDD; hence, significant difference was determined between the two groups ($p<0.01$).

Figure 4. The percentage of vitamin D deficiency (VDD) according to obesity in osteoarthritis (OA) and control group. There was no significant difference in the percentage of VDD in the OA and control groups in obesity patients. In non-obesity patients, there was significant difference between the two groups.




VDD according to season of sampling

Table 2 shows the percentage of vitamin D deficiency, insufficiency, and sufficiency by season of sampling. Between the two groups, there was a significant difference in the prevalence of VDD according to the seasons in summer, autumn, and winter, and the difference was the largest in summer (24.1% points, $p<0.001$), followed by winter (15.4% points, $p<0.001$), and autumn (13.3% points, $p=0.03$) (Figure 5). Seasonal differences in VDD prevalence in the OA group were significant only in winter (81.2%) and autumn (70.1%) ($p=0.04$). In the control group, there were significant differences between summer and spring ($p<0.001$) and winter and summer ($p<0.001$).

Table 2. Vitamin D status according to season of sampling in osteoarthritis (OA) and control group

| Group | Season of sampling | Vitamin D | Vitamin D | Vitamin D | Total |
|---------|--------------------|--------------|---------------|-------------|-------|
| | | deficiency | insufficiency | sufficiency | |
| | | <i>n</i> (%) | | | |
| OA | Spring | 93 (74.4) | 22 (17.6) | 10 (8.0) | 125 |
| | Summer | 95 (72.5) | 26 (19.8) | 10 (7.6) | 131 |
| | Autumn | 75 (70.1) | 23 (21.5) | 9 (8.4) | 107 |
| | Winter | 112 (81.2) | 21 (15.2) | 5 (3.6) | 138 |
| Control | Spring | 124 (66.6) | 40 (21.3) | 24 (12.8) | 188 |
| | Summer | 88 (48.4) | 56 (30.8) | 38 (20.9) | 182 |
| | Autumn | 88 (56.8) | 31 (26.5) | 26 (16.8) | 155 |
| | Winter | 129 (65.8) | 40 (20.4) | 27 (13.8) | 196 |

Figure 5. The percentage of vitamin D deficiency according to the season of sampling in the osteoarthritis (OA) and control group. Between the two groups, a significant difference was observed in the percentage of VDD in summer, autumn, and winter (asterisk). Seasonal differences in VDD prevalence in the OA group were significant only in winter and autumn. In the control group, there were significant differences in VDD prevalence between summer and spring and between winter and summer.



Discussion

This study had the largest cohort of patients undergoing TKA who were screened for VDD and compared with a matched healthy population. A high VDD prevalence was evident in patients who underwent TKA, with 75.0% of the OA group having 25(OH)D levels of <20 ng/dL. Moreover, a significant difference was observed between the two groups.

Heidari et al.¹⁸ determined the status of serum 25(OH)D levels in patients with knee OA compared with controls and have reported that in the entire population, the mean serum 25(OH)D level in patients with OA was not significantly lower than that of the controls. However, this study showed that the serum 25(OH)D level of the case group who had knee OA and underwent TKA was lower than that of the control group, and in that the average age of the total population in this study was 67.3 years, this study showed a different result from the previous study that the serum 25(OH)D level was low only in patients <60 years of age among knee OA patients.¹⁸ Bhattoa et al.¹⁹ characterized the prevalence of VDD and vitamin D insufficiency in healthy men aged >50 years and have reported that the prevalence of hypovitaminosis D was 52.9%. As a result of analyzing only men >55 years of age in the control group of this study, the prevalence of hypovitaminosis D among 185 men with an average age of 68.4 was 75.7% (VDD, 47.6%; vitamin D insufficiency, 28.1%), which was higher than that of the previous study.

Arima et al.²⁰ have reviewed several epidemiological studies, which have assessed the correlation between serum 25(OH)D levels and OA in elderly individuals and have reported that data on the association between serum 25(OH)D levels and OA are contrasting. Although this study did not directly compare the OA grades of the two groups, TKA was generally performed in patients with a moderate or advanced grade of knee OA, and even with the same OA grade, the patients who had more severe pain underwent TKA. Therefore, this study's results can be interpreted as the serum 25(OH)D level having an inverse association with the grade of knee OA, or the severity of pain caused by knee OA. The authors speculated that the reason for the decrease in serum 25(OH)D level in knee OA patients was that the overall number of activities including outdoor activities decreased due to knee pain and this is consistent with a previous study²¹ that has evaluated the association between serum 25(OH)D level and physical activity. Klingberg et al.²¹ explored parameters associated with 25(OH)D in a healthy adult population and serum 25(OH)D was significantly higher in the physically active population who performed physical exercise regularly at least once a week.

Goula et al.¹² have reported a significant positive association between serum 25(OH)D levels and the male sex in patients who were scheduled for knee or hip joint replacement. The result of a previous study can be interpreted

as the reason that women have less outdoor activities than men.²⁰ However, in the present study, no significant difference was noted between men and women in the OA group ($p=0.06$). The authors believed that the reason of the significantly higher VDD percentage for both men and women in the OA group compared to the matched control group was that patients in the OA group had severe knee pain enough to undergo TKA, and it is possible that the knee pain affected the outdoor activity level or sun exposure. Similarly, the authors believed that the reason why there was no significant difference in VDD percentage between men and women in the OA group was that patients with moderate or advanced OA had similar degrees of restriction of movement and decrease in outdoor activities due to similar severe knee pain.

Maier et al.¹¹ assessed risk factors for insufficient 25(OH)D levels in patients aged >70 years and admitted to an orthopedic surgery department and have reported that patients presenting with obesity were more likely to have low 25(OH)D levels. In this study, after matching between the patients in the OA group and the control group, the standardized mean difference of BMI between the two values was <0.2; however, the serum 25(OH)D level in the OA group was significantly lower than that of the control group. A significant difference was observed in BMI values by vitamin D status in the total population and control groups ($p<0.01$, 0.011); however, in the OA group, the difference in BMI values by vitamin D status was not significant ($p=0.63$). In the analysis of non-obesity patients, the VDD percentage of the OA group (74.7%) was significantly higher than that of the control group (58.7%). Regarding these results, the authors hypothesized that in the OA group, not only the association between serum 25(OH)D level and obesity but also a decrease in outdoor activity level and in sun exposure due to knee OA were added.

In general, VDD is known to decrease in summer,²¹ and in this study, it was confirmed that healthy participants in the control group significantly decreased VDD in summer compared to spring and winter. However, VDD was confirmed in more than 70% of OA patients who underwent TKA regardless of the season, and in winter, >80% of patients had VDD. Regarding these results, the authors thought that it was because the outdoor activity level or sun exposure did not change according to the season in patients with severe knee pain enough to undergo TKA. The authors thought that the greatest difference (24% points) between the two groups appeared in summer because of the characteristics of the OA group, where VDD was high regardless of the season, and the general population, where VDD decreased in summer.

The strength of this study was that healthy participants whose age, sex, weight, BMI, and chemistry were matched were used to compare serum 25(OH)D level in patients who underwent TKA. However, this study had several limitations. First, we did not analyze whether the knee X-rays were taken or OA on X-rays was graded in

the control group. However, for the reasons mentioned above, it can be estimated that the healthy participants in the control group had an OA grade less than K-L grade II or were not painful enough to receive TKA even if their OA grade on X-ray was higher than K-L grade III. Second, markers of bone metabolism, including serum parathyroid hormone were not analyzed. However, no difference in serum calcium and phosphorus levels between groups may represent similar calcium homeostasis in both groups. Third, 40% of patients in the case group were excluded after PSM. However, this was excluded during the process of matching the two groups. After matching, no demographic difference was noted between the two groups, and even after matching, a large population was still analyzed. Fourth, clinical outcomes could not be evaluated in the case group because this study compared case group that underwent TKA with a healthy control group. Fifth, a more severe knee OA had a lower bone mineral density (BMD) and serum 25(OH)D level;²³ hence, it is also a limitation that BMD was not used as a matching variable in this study. Finally, because this study retrospectively analyzed data that was collected and organized only for specific items selected in advance, the levels of outdoor physical activity or sun exposure, well known as significant determinants of vitamin D status, could not be collected.

Conclusion

This study showed that the serum 25(OH)D levels in the OA group was 15.3 ng/mL, which was significantly lower than that in the control group, and the prevalence of VDD in the OA group was 75.0%, which was significantly higher than that in the control group (59.4%). In the OA group, no differences were observed between males and females and according to obesity status. Additionally, patients in the OA group had more severe VDD in summer than healthy participants, and VDD was confirmed in more than 80% of patients in the OA group in winter.

1 **References**

- 2 1. Garfinkel RJ, Dilisio MF, Agrawal DK (2017) Vitamin D and Its Effects on Articular Cartilage and
3 Osteoarthritis. *Orthop J Sports Med* 5:2325967117711376. doi:10.1177/2325967117711376
- 4 2. Tomlinson PB, Joseph C, Angioi M (2015) Effects of vitamin D supplementation on upper and lower body
5 muscle strength levels in healthy individuals. A systematic review with meta-analysis. *J Sci Med Sport* 18:575-
6 580. doi:10.1016/j.jsams.2014.07.022
- 7 3. Dhaliwal R, Aloia JF (2017) Effect of Vitamin D on Falls and Physical Performance. *Endocrinol Metab Clin*
8 North Am 46:919-933. doi:10.1016/j.ecl.2017.07.004
- 9 4. Barker T, Henriksen VT, Martins TB, Hill HR, Kjeldsberg CR, Schneider ED, Dixon BM, Weaver LK (2013)
10 Higher serum 25-hydroxyvitamin D concentrations associate with a faster recovery of skeletal muscle strength
11 after muscular injury. *Nutrients* 5:1253-1275. doi:10.3390/nu5041253
- 12 5. Oh JH, Kim SH, Kim JH, Shin YH, Yoon JP, Oh CH (2009) The level of vitamin D in the serum correlates with
13 fatty degeneration of the muscles of the rotator cuff. *J Bone Joint Surg Br* 91:1587-1593. doi:10.1302/0301-
14 620x.91b12.22481
- 15 6. Maier GS, Maus U, Lazovic D, Horas K, Roth KE, Kurth AA (2016) Is there an association between low serum
16 25-OH-D levels and the length of hospital stay in orthopaedic patients after arthroplasty? *J Orthop Traumatol*
17 17:297-302. doi:10.1007/s10195-016-0414-y
- 18 7. Shin KY, Park KK, Moon SH, Yang IH, Choi HJ, Lee WS (2017) Vitamin D deficiency adversely affects early
19 post-operative functional outcomes after total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 25:3424-
20 3430. doi:10.1007/s00167-016-4209-8
- 21 8. Jansen JA, Haddad FS (2013) High prevalence of vitamin D deficiency in elderly patients with advanced
22 osteoarthritis scheduled for total knee replacement associated with poorer preoperative functional state. *Ann R*
23 *Coll Surg Engl* 95:569-572. doi:10.1308/003588413x13781990150374
- 24 9. Maniar RN, Patil AM, Maniar AR, Gangaraju B, Singh J (2016) Effect of Preoperative Vitamin D Levels on
25 Functional Performance after Total Knee Arthroplasty. *Clin Orthop Surg* 8:153-156.
26 doi:10.4055/cios.2016.8.2.153
- 27 10. Piuzzi NS, George J, Khlopas A, Klika AK, Mont MA, Muschler GF, Higuera CA (2018) High prevalence and
28 seasonal variation of hypovitaminosis D in patients scheduled for lower extremity total joint arthroplasty. *Ann*
29 *Transl Med* 6:321. doi:10.21037/atm.2018.08.21
- 30 11. Maier GS, Seeger JB, Horas K, Roth KE, Kurth AA, Maus U (2015) The prevalence of vitamin D deficiency
31 in patients with vertebral fragility fractures. *Bone Joint J* 97:89-93. doi:10.1302/0301-620x.97b1.34558

- 32 12. Goula T, Kouskoulis A, Drosos G, Tselepis AS, Ververidis A, Valkanis C, Zisimopoulos A, Kazakos K (2015)
33 Vitamin D status in patients with knee or hip osteoarthritis in a Mediterranean country. *J Orthop Traumatol* 16:35-
34 39. doi:10.1007/s10195-014-0322-y
- 35 13. Ginde AA, Liu MC, Camargo CA Jr. (2008) Demographic differences and trends of vitamin D insufficiency
36 in the US population, 1988-2004. *Arch Intern Med.* 169:626-632. doi: 10.1001/archinternmed.2008.604.
- 37 14. Choi HS, Oh HJ, Choi H, Choi WH, Kim JG, Kim KM, Kim KJ, Rhee Y, Lim SK. (2011) Vitamin D
38 insufficiency in Korea--a greater threat to younger generation: the Korea National Health and Nutrition
39 Examination Survey (KNHANES) 2008. *J Clin Endocrinol Metab* 96:643-651. doi: 10.1210/jc.2010-2133.
- 40 16. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, Murad MH, Weaver CM
41 (2011) Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice
42 guideline. *J Clin Endocrinol Metab* 96:1911-1930. doi:10.1210/jc.2011-0385
- 43 15. Kellgren JH, Lawrence JS (1957) Radiological assessment of osteo-arthrosis. *Annals of the rheumatic diseases*
44 16:494-502. <https://doi.org/10.1136/ard.16.4.494>
- 45 17. Apovian CM (2016) Obesity: definition, comorbidities, causes, and burden. *Am J Manag Care* 22:176-185.
- 46 18. Heidari B, Heidari P, Hajian-Tilaki K (2011) Association between serum vitamin D deficiency and knee
47 osteoarthritis. *Int Orthop* 35:1627-1631. doi:10.1007/s00264-010-1186-2
- 48 19. Bhattoa HP, Nagy E, More C, Kappelmayer J, Balogh A, Kalina E, Antal-Szalmas P (2013) Prevalence and
49 seasonal variation of hypovitaminosis D and its relationship to bone metabolism in healthy Hungarian men over
50 50 years of age: the HunMen Study. *Osteoporos Int* 24:179-186. doi:10.1007/s00198-012-1920-2
- 51 20. Arima K, Mizukami S, Nishimura T, Tomita Y, Nakashima H, Abe Y, Aoyagi K (2020) Epidemiology of the
52 association between serum 25-hydroxyvitamin D levels and musculoskeletal conditions among elderly individuals:
53 a literature review. *J Physiol Anthropol* 39:38. doi:10.1186/s40101-020-00249-3
- 54 21. Klingberg E, Oleröd G, Konar J, Petzold M, Hammarsten O (2015) Seasonal variations in serum 25-hydroxy
55 vitamin D levels in a Swedish cohort. *Endocrine* 49:800-808. doi: 10.1007/s12020-015-0548-3.
- 56 22. Kim SH, Oh JE, Song DW, Cho CY, Hong SH, Cho YJ, Yoo BW, Shin KS, Joe H, Shin HS, Son DY (2018)
57 The factors associated with Vitamin D deficiency in community dwelling elderly in Korea. *Nutr Res Pract* 12:387-
58 395. doi: 10.4162/nrp.2018.12.5.387.
- 59 23. Linde KN, Puhakka KB, Langdahl BL, Søballe K, Krog-Mikkelsen I, Madsen F, Stilling M (2017) Bone
60 Mineral Density is Lower in Patients with Severe Knee Osteoarthritis and Attrition. *Calcif Tissue Int* 101:593-
61 601. doi:10.1007/s00223-017-0315-y

요약 (국문)

목적: 본 연구에서는 매칭 분석을 통해 슬관절 인공관절 전치환술을 받은 환자와 건강한 참가자 간의 혈청 25-OH 비타민 D 수치와 비타민 D 결핍의 유병률을 조사하고 비교하였다.

대상 및 방법: 전체적으로 슬관절 인공관절 전치환술을 받은 824명의 환자가 연구 군에 포함되었고 대조 군은 동일 의료 기관에서 건강 검진을 받은 2,794명의 참가자로 구성되었다. 성별, 나이, 체중, 체질량지수, 혈액 화학 검사, 채혈 일자에 따라 성향 점수 매칭을 통해 대조 군을 조정하였다. 성향 점수 매칭을 통해 연구 군과 대조 군에 각각 501명, 721명이 배정되었다. 환자의 체질량 지수가 30kg/m²인 경우를 비만으로 분류하였다. 혈청 25-OH 비타민 D를 포함한 혈액 화학 수준을 조사하였으며, 비타민 D 결핍은 <20 ng/mL로 정의하였다.

결과: 매칭 후, 두 군의 혈청 25-OH 비타민 D 수치는 각각 15.3 ng/mL와 19.9 ng/mL이었고, 두 군의 비타민 D 결핍 유병률은 각각 75.0%와 59.4%였으며, 통계학적으로 유의한 차이가 있었다. 연구 군에서 성별과 비만 상태에 따른 차이는 관찰되지 않았다. 연구 군에서 계절별 비타민 D 결핍은 가을과 겨울에 채혈한 환자간 유의한 차이가 관찰되었다. 두 군에서 계절별 비타민 D 결핍의 차이는 여름에 가장 큰 폭의 차이를 보였다 (24.1% 포인트).

결론: 슬관절 인공관절 전치환술을 받은 환자는 성향 점수로 매칭된 건강한 대조 군과 비교하여 혈청 25-OH 비타민 D 수치는 낮고 비타민 D 결핍 비율은 높았다. 슬관절 인공관절 전치환술을 받은 환자에서 성별이나 비만 여부에 따른 비타민 D 결핍율에는 차이가 없었으며, 비타민 D 결핍율은 4계절 모두 70% 이상으로 확인되었다. 따라서, 슬관절 인공관절 전치환술을 받은 환자에서 성별, 비만, 계절과 무관하게 비타민 D 결핍에 대한 포괄적인 인지가 필요하겠다.

근거 수준: 후향적 코호트 연구; Level of evidence III

중심 단어: 25-OH 비타민 D, 슬관절, 골관절염, 인공관절 전치환술, 건강 검진, 비타민 D 결핍